



## **Matrix metalloprotease-sensitive doxorubicin-loaded liposomes for enhanced anticancer activity.**

**Clergeaud Veiga, Gael; Fliedner, Frederikke P.; Hansen, Anders Elias; Melander, Fredrik; Kjær, Andreas; Andresen, Thomas Lars**

*Publication date:*  
2016

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Clergeaud Veiga, G., Fliedner, F. P., Hansen, A. E., Melander, F., Kjær, A., & Andresen, T. L. (2016). *Matrix metalloprotease-sensitive doxorubicin-loaded liposomes for enhanced anticancer activity..* Poster session presented at The 43rd Annual Meeting & Exposition of the Controlled Release Society, Seattle, WA, United States.

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.




# Matrix Metalloprotease-Sensitive Doxorubicin-Loaded Liposomes for Enhanced Anticancer Activity

Gael Clergeaud<sup>1,2</sup>, Frederikke P. Fliedner<sup>3</sup>, Anders E. Hansen<sup>1,2,3</sup>, Fredrik Melander<sup>1,2</sup>, Andreas Kjær<sup>3</sup>, Thomas L. Andresen<sup>1,2</sup>

<sup>1</sup>Department of Micro- and Nanotechnology, Technical University of Denmark, Building 423, DK-2800 Kgs. Lyngby, Denmark; <sup>2</sup>Centre for Nanomedicine and Theranostics, Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark; <sup>3</sup>Department of Clinical Physiology, Nuclear Medicine & PET, and Cluster for Molecular Imaging, Rigshospitalet, Copenhagen University Hospital and Faculty of Health Sciences, University of Copenhagen, Blegdamsvej 3B, DK-2200 Copenhagen, Denmark.

## Purpose


Long-circulating PEGylated liposomal formulations of doxorubicin (DOX) have shown to increase drug accumulation in tumors and reduce dose-limiting effects of DOX (e.g. cardiotoxicity); however these systems have only achieved a moderate improvement in anti-tumor activity due to poor cellular internalization and drug bio-availability<sup>1</sup>. We have recently developed an enzyme-sensitive liposome system that exploits the proteolytic action of matrix metalloproteases (MMPs), over-expressed in broad range of cancers, to produce the detachment of the PEG-coat and induce a membrane charge shift that will promote their internalization by tumor cells and the intracellular drug release<sup>2</sup>. We aim to demonstrate that DOX formulated into cationic liposomes covered with a detachable negatively-charged PEG-coating are efficient nanocarriers to enhance the delivery of DOX to tumor cells and improve the anti-tumor activity.




## Results

The MMP-sensitive formulation increased the uptake of the liposomes by the cells upon proteolytic activation, resulting in an increased intracellular delivery of DOX and improved cytotoxic activity as compared with DOXIL (Fig. 1-3). The DOX-loaded MMP-sensitive liposomes showed a reduced blood circulation profile as compared with DOXIL (Fig. 4), suggesting a premature leakage of DOX from the carrier and/or fast removal by the RES. Thus, DOX levels found in the tumor were reduced whereas in the spleen were significantly higher. Encapsulation of DOX within MMP-sensitive liposomes resulted in slightly better tumor growth inhibition compared to the Free DOX but failed to provide similar therapeutic efficacy as DOXIL.

### *In vitro* cell uptake and growth inhibition of DOX formulated in MMP-sensitive liposomes



### *In vivo* evaluation of DOX-loaded MMP-sensitive liposomes



## Conclusions

The results show the remarkable capacity of the MMP-triggered liposomes to increase the cellular internalization of DOX and efficiently improve the *in vitro* growth inhibition of cancer cells compared to DOXIL. However, the pharmacokinetic and biodistribution profiles demonstrated poor circulating properties and tumor accumulation, leading to an unimproved *in vivo* therapeutic potential compared to DOXIL. Further investigation on designing a more stable MMP-sensitive formulation should be carried on in order to ensure long circulating properties of the drug.

## References

- Barenholz Y. J. *Control. Release*. 2012 (160) 117-134.
- Gjetting T., Jølcck R., Andresen T. L. *Adv. Healthc. Mater.* 2014 (3) 1107-1118.
- Laginha K. M., Verwoert S., Charrois G. J. R., Allen T. M. *Clin. Cancer Res.* 2005 (19) 6944-6949.

gaelc@nanotech.dtu.dk

